

TREE SPECIES (SCIENTIFIC NAME)

Pouteria bilocularis

PERU COMMON NAME

Caimitillo

TREE FAMILY

SAPOTACEAE

AVERAGE LEAF SIZE (CM)

11.86cm × **6.04cm**
Length Width

DISTRIBUTION



NATIVE TO PERU

ELEVATIONAL RANGE (M)

350–1500M

TREE HEIGHT

MEDIUM (20–35M)



NATIVE TO

Region: Americas

Latin America: Bolivia, Brazil, Colombia, Ecuador, Guyana, Peru, Venezuela

COFFEE AGROFORESTRY INFORMATION

COFFEE SYSTEM



ARABICA

COFFEE IMPACT

Unknown

TREE MANAGEMENT

86% germination rates were obtained in Brazil after seeds were extracted from fruit, washed for 30 minutes, dried in shade for 18 hours at ambient temperatures, and planted at 1 in depth in substrate of sand and sawdust.

CULTIVATION



PLANTED

PREVALENCE

Not Common in Coffee Agroforestry

TREE BENEFITS AND USES

FARMER USES

Unknown

FARM SERVICES

Unknown

BIODIVERSITY BENEFITS

✓ **YES**

The fruits are eaten by the monkey *Cebus apella*.

Last Updated: August 15, 2023

Cruz, E. D. (2005). Quantitative characteristics of fruits and seeds of *Pouteria pachycarpa* Pires-Sapotaceae. *Revista Brasileira de Sementes*, 27, 159-164.;

Pennington, T.D. & R. Bernal 2022-7-12. *Pouteria bilocularis* (H.J.P.Winkl.) Baehni En Bernal, R., S.R. Gradstein & M. Celis (eds.). 2015.;

Catálogo de plantas y líquenes de Colombia. Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá. <http://catalogoplantasdecolombia.unal.edu.co>;

WFO (2022): *Pouteria bilocularis* (H.J.P.Winkl.) Baehni. Accessed on: 12 Jul 2022. Published on the Internet <http://www.worldfloraonline.org/taxon/wfo-0000281573>;

Mendoza, A. L. M., & Guerrero, M. H. (2010). Catálogo de los árboles y afines de la Selva Central del Perú. *Amaldoa*, 17, 203-242.